

****TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

The Peculiar Cooling Flow Cluster RX J0820.9+0752

C. M. Bayer-Kim, C. S. Crawford, S. W. Allen, A. C. Fabian

*University of Cambridge, Institute of Astronomy, Madingley Road,
Cambridge, CB3 0HA, UK*

A. C. Edge

*Department of Physics, University of Durham, South Road, Durham
DH1 3LE*

Abstract. This poster is a summary of a paper on the subject (Bayer-Kim et al. 2002) that has been submitted for publication in MNRAS. We present X-ray, $H\alpha$, radio, and optical data, both imaging and spectroscopy, of the cluster RX J0820.9+0752 and its peculiar central cluster galaxy (CCG). We announce the discovery of several isolated off-nuclear patches of blue light with strong line-emission (total $L(H\alpha) \sim 10^{42} \text{ergs}^{-1}$) and distinct ionizational and kinematic properties, embedded in a region of $H\alpha$ /X-ray emission. We propose and investigate a scenario in which a secondary galaxy also featured in our observations has moved through a cooling wake produced by the CCG, thereby producing the observed clumpy morphology and helping to trigger star-formation within the clumps.

1. RX J0820.9+0752 At Different Wavelengths

From the strongly peaked profile in our 9.4 ks *Chandra* observation (Fig. 1a), we derive that the hot intracluster gas in the cluster RX J0820.9+0752 is cooling within a radius of $r \approx 20$ kpc. The mass deposition rate of a few tens of solar masses per year is consistent with the new, reduced cooling flow scenario proposed by Voigt et al. (2002) as a possible solution to the problem of the missing soft X-ray luminosity in cooling flow clusters (Peterson et al. 2001). The X-ray emission appears extended to the NW and is coincident with a luminous $H\alpha$ nebula seen in an AAT image taken through a narrow-band filter in the $H\alpha$ light (Fig. 1b). The cluster contains only a weak radio source (flux density of 0.45 ± 0.13 mJy at 4.89 GHz as determined from a complete VLA snapshot campaign; Edge et al. 2002, in prep.), but in an HST image its central galaxy shows highly peculiar morphology with two arcs of clumped emission to the NE and a secondary elliptical to the SE, opposite the X-ray/ $H\alpha$ feature (Fig. 1c). Several other isolated blobs are also apparent scattered further out to the NW and E.




Figure 1. (a) The slightly smoothed 0.5–2 keV *Chandra* image of the cluster emission from RX J0820.9+0752 with the optical contours overlaid (top). (b) Continuum-subtracted $H\alpha$ emission around the central galaxy. The image shown is 20.5×16 arcsec (middle). (c) F606W HST image ($\Delta\lambda \approx 4490 - 5910\text{\AA}$ at the redshift of the object) of the central region of the cluster on the same scale as the $H\alpha$ image (bottom). North is to the top and East to the left in all the figures.





Figure 2. (a) Plot of $[\text{NII}]/\text{H}\alpha$ as a function of radius. The positions of three blue blobs are marked. Note their association with lower values of $[\text{NII}]/\text{H}\alpha$ (top). (b) FWHM plotted vs r . The same blobs as above are denoted. Again they show deviations from the general trend (bottom).

2. Optical Spectroscopy

We acquired optical spectra on the red (3300 – 5000Å) and blue (5100 – 6800Å rest wavelength) arms of ISIS (WHT, La Palma) to cover both the known blue excess and the strong red line emission in this object. The high spectral resolution allowed us to extract separate spectra of some of the blobs seen in the HST image. They are regions of very strong line emission ($\text{H}\alpha$ slit Flux $\sim 10^{-15} \text{erg s}^{-1} \text{cm}^{-2} \text{\AA}^{-1}$), associated with a dust-rich environment ($E(B - V)$ varies between about 0.3 – 0.7 as determined from the Balmer decrement). Diagnostic line-ratios sensitive to ionization properties show considerable variation throughout the whole object, but with consistent deviations associated with the blobs (Fig. 2a). The blobs are also kinematically distinct regions with separate v_{rad} and FWHM properties (Fig. 2b).

We find the blobs to be associated with very blue continua, whereas the nuclei of the two galaxies are perfectly consistent with those of “normal” ellipticals. We have fit the blue blob spectra using models constructed from an average template CCG (Crawford et al. 1999) and a variety of MS stellar spectra from Pickles’ (1998) stellar spectral flux library (Fig. 3). It is possible to explain the blue excess by the presence of significant numbers of early stars (up to $\sim 10^5$ O-stars) from recent star-bursts of varying ages.

Figure 3. Fit to one of the blue blobs. The spectrum is represented by the dashed line, the model by the solid one. The dotted spectrum is the difference between the late component of the model and the blob spectrum, clearly showing the blue excess and the Balmer and [OII] λ 3727Å emission lines.



3. Conclusions

The association of the H α nebula with the X-ray emission is reminiscent of a similar feature found recently in A1795 (Fabian et al. 2001b). The most probable explanation given in this paper is that the filament represents a “cooling wake”, produced by the central cluster galaxy moving through the hot ICM. We show that the same explanation can be applied to RX J0820.9+0752, where the weakness of the radio source practically rules out the possibility that it has played a significant role in producing the wake. We find that the energy output through the strong line-emission is consistent with ionization from energetic photons produced by young stars. However, we observe a velocity offset of up to 200 km s $^{-1}$ between the young stellar components and the line-emitting gas in the filament, suggesting they are not co-spatial. This could indicate that massive stars are not the only ionization source for the nebula. Maybe processes such as the “cold mixing model” proposed by Fabian et al. (2001a) can account for at least part of the observed line-emission.

An important factor in producing the clumpy morphology and triggering the star-bursts seems to be the secondary elliptical galaxy, whose velocity properties relate it to the cooling gas. We propose a scenario in which the secondary galaxy passes through the cooling wake produced by the CCG.

References

- Bayer-Kim, C. M., Crawford, C. S., Allen, S. W., Edge, A. C., Fabian, A. C. 2002, MNRAS, submitted
- Fabian, A. C., Sanders, J. S., Ettori, S., Taylor, G. B., Allen, S. W., Crawford, C. S., Iwasawa, K., & Johnstone, R. M. 2001a, MNRAS, 321, L20
- Fabian, A. C., Mushotzky, R. F., Nulsen, P. E. J., & Peterson, J. R. 2001b, MNRAS, 321, L33

- Crawford C. S., Allen, S. W., Ebeling, H., Edge, A. C., & Fabian, A. C. 1999, MNRAS, 306, 857
- Peterson, J. R., Paerels, F. B. S., Kaastra, J. S., Arnaud, M., Reiprich, T. H., Fabian, A. C., Mushotzky, R. F., Jernigan, J. G., & Sakelliou, I. 2001, A&A, 365, 104
- Pickles A. J. 1998, PASP, 110, 863
- Voigt, L. M., Schmidt, R. W., Fabian, A. C., Allen, S. W., & Johnstone, R. M. 2002, MNRAS, submitted (astro-ph/0203312)